

ALBON® ORAL SUSPENSION - 5%

Rx

Pfizer Animal Health

Sulfadimethoxine

(Brand of Sulfadimethoxine)

Each teaspoonful (5 mL) contains 250 mg sulfadimethoxine.


Dogs and Cats - To be used in the treatment sulfadimethoxine-sensitive bacterial infections in dogs and cats and bacterial enteritis associated with coccidiosis in dogs.

Not for Human Use

CAUTION: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION: Albon is a low-dosage, rapidly absorbed, long-acting sulfonamide, effective for the treatment of a wide range of bacterial infections commonly encountered in dogs and cats.

Sulfadimethoxine is a white, almost tasteless and odorless compound. Chemically, it is N¹-(2,6-dimethoxy-4-pyrimidinyl) sulfanilamide. The structural formula is:



ACTIONS: Sulfadimethoxine has been demonstrated clinically or in the laboratory to be effective against a variety of organisms, such as streptococci, klebsiella, proteus, shigella, staphylococci, escherichia, and salmonella.^{1,2} These organisms have been demonstrated in respiratory, genitourinary, enteric, and soft tissue infections of dogs and cats.

The systemic sulfonamides which include sulfadimethoxine are bacteriostatic agents. Sulfonamides competitively inhibit bacterial synthesis of folic acid (pteroylglutamic acid) from para-aminobenzoic acid. Mammalian cells are capable of utilizing folic acid in the presence of sulfonamides.

The tissue distribution of sulfadimethoxine, as with all sulfonamides, is a function of plasma levels, degree of plasma protein binding, and subsequent passive distribution in the tissues of the lipid-soluble unionized form. The relative amounts are determined by both its pKa and by the pH of each tissue. Therefore, levels tend to be higher in less acid tissue and body fluids or those diseased tissues having high concentrations of leucocytes.²

In the dog, sulfadimethoxine is not acetylated as in most other animals, and it is excreted predominantly as the unchanged drug.³ Sulfadimethoxine has a relatively high solubility at the pH normally occurring in the kidney, precluding the possibility of precipitation and crystalluria. Slow renal excretion results from a high degree of tubular reabsorption,⁴ and plasma protein binding is very high, providing a blood reservoir of the drug. Thus, sulfadimethoxine maintains higher blood levels than most other long-acting sulfonamides. Single, comparatively low doses of Albon give rapid and sustained therapeutic blood levels.¹

To assure successful sulfonamide therapy (1) the drug must be given early in the course of the disease, and it must produce a high sulfonamide level in the body rapidly after administration, (2) therapeutically effective sulfonamide levels must be maintained in the body throughout the treatment period, (3) treatment should continue for a short period of time after the clinical signs have disappeared, and (4) the causative organisms must be sensitive to this class of drugs.

TOXICITY AND SAFETY: Data regarding acute and chronic toxicities of sulfadimethoxine indicate the drug is very safe. The LD₅₀ in mice is greater than 2 g/kg of body weight when administered intraperitoneally and greater than 16 g/kg when administered orally. In dogs receiving massive single oral doses of 3.2 g/kg of body weight, diarrhea was the only adverse effect observed. Dogs given 160 mg/kg of body weight orally daily for 13 weeks showed no signs of toxicity.

INDICATIONS: Albon is indicated for the treatment of a wide range of respiratory, genitourinary tract, enteric, and soft tissue infections. For example: tonsillitis, pustular dermatitis, pharyngitis, anal gland infections, bronchitis, abscesses, pneumonia, wound infections, cystitis, bacterial enteritis, nephritis, canine salmonellosis, metritis, bacterial enteritis associated with coccidiosis in dogs, pyometra when caused by streptococci, staphylococci, escherichia, salmonella, klebsiella, proteus or shigella organisms, sensitive to sulfadimethoxine.

LIMITATIONS: Sulfadimethoxine is not effective in viral or rickettsial infections, and as with any antibacterial agent, occasional failures in therapy may occur due to resistant microorganisms. The usual precautions in sulfonamide therapy should be observed.

PRECAUTION: During treatment period, make certain that animals maintain adequate water intake.

If animals show no improvement within 2 or 3 days, reevaluate your diagnosis.

DOSAGE AND ADMINISTRATION:

Initial dose: 25 mg/lb (55 mg/kg) of animal body weight.

Subsequent Daily Doses: 12.5 mg/lb (27.5 mg/kg) of animal body weight.

Albon Oral Suspension 5%

Dogs and cats should receive 1 teaspoonful of Albon Oral Suspension 5% per 10 lb of body weight (25 mg/lb or 55 mg/kg) as an initial dose, followed by ½ teaspoonful per 10 lb of body weight (12.5 mg/lb or 27.5 mg/kg) every 24 hours thereafter. Representative weights and doses are indicated in the following table:

Animal Weight	Initial Dose 25 mg/lb (55 mg/kg)	Subsequent Daily Doses 12.5 mg/lb (27.5 mg/kg)
5 lb (2.2 kg)	½ tsp (2½ mL)	¼ tsp (1¼ mL)
10 lb (4.5 kg)	1 tsp (5 mL)	½ tsp (2½ mL)
20 lb (9.1 kg)	2 tsp (10 mL)	1 tsp (5 mL)
40 lb (18.2 kg)	4 tsp (20 mL)	2 tsp (10 mL)
80 lb (36.4 kg)	8 tsp (40 mL)	4 tsp (20 mL)

Treatment may be initiated with Albon Injection 40% to obtain effective blood levels almost immediately or to facilitate treatment of the fractious animal.

Length of treatment depends on the clinical response. In most cases treatment for 3-5 days is adequate. Treatment should be continued until the animal is asymptomatic for 48 hours.

HOW SUPPLIED: Albon is available in the following dosage forms for dogs and cats:

Albon Tablets: 125 mg, 250 mg, or 500 mg sulfadimethoxine per tablet.

Albon Oral Suspension 5%: 2- and 16-oz bottles; each tsp (5 mL) contains 250 mg sulfadimethoxine in a custard-flavored carrier.

Albon Injection 40%: 100-mL multiple-dose vials.

Each mL contains 400 mg sulfadimethoxine compounded with 20% propylene glycol, 1% benzyl alcohol, 0.1 mg disodium edetate, 1 mg sodium formaldehyde sulfoxylate, and pH adjusted with sodium hydroxide.

REFERENCES:

1. Data on file from Hoffmann-La Roche Inc., Nutley, New Jersey.
2. Stowe CM: The sulfonamides. In Jones LM (ed), *Veterinary Pharmacology and Therapeutics*, Ames, Iowa, Iowa State University Press, chapter 33, 1965.
3. Bridges JW, Kirby MR, Walker SR, *et al*: Species differences in the metabolism of sulfadimethoxine. *Biochem J* 109:851,1968.
4. Baggot JD: Some aspects of drug persistence in domestic animals. *Res Vet Sci* 11 (2):130, 1970.

NADA #015-102, Approved by FDA for Tablets

NADA #043-785, Approved by FDA for Suspension

Manufactured by:

Roche Vitamins and Fine Chemicals

A division of Hoffmann-La Roche Inc.

Nutley, NJ 07110, USA

75-8430-01

February 1994

Disclaimer: Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the product label or package insert. Compendium Code No.: 03691006